

Paradoxes in dermatology

Keshavmurthy A. Adya, Arun C. Inamadar, Aparna Palit

Department of
Dermatology,
Venereology and Leprosy,
Shri B. M. Patil Medical
College, Hospital and
Research Center, Bijapur
Liberal Development
Education University,
Bijapur, Karnataka, India

ABSTRACT

Many paradoxical phenomena related to clinical, immunological, and therapeutic dermatology have been described. While some of them can be explained logically, the cause for others can only be speculated. Whenever encountered in clinical practice, background knowledge of such paradoxes may be useful to the clinician.

Key words: Granuloma paradox, paraben paradox, paradoxical hypertrichosis

Diagnosing dermatological conditions has always been intriguing. Many unrelated conditions present with apparently similar morphology that often prove challenging to the untrained eyes. Furthermore, the existence of several complicated synonyms for apparently ordinary conditions do not help matters either. To add to this, several paradoxes and misnomers related to clinical, immunological, and therapeutic dermatology have also been described. A basic knowledge of such paradoxes, in addition to the usual, may be helpful when such situations are encountered in clinical practice.

A “paradox” is defined as a “*seemingly absurd or contradictory statement or proposition which when investigated may prove to be well founded or true.*”^[1] In contrast, a “misnomer” is a completely inaccurate statement or assertion. There are some well-known paradoxes in dermatology, some of which can be explained logically or based on evidence, while for some, only assumptions or postulates are put forth. Furthermore, some of these paradoxes are yet to be ascertained any explanations. Following are some of such paradoxes in dermatology compiled under clinical, immunological, dermatopathological, and therapeutic categories.

PARADOXES IN CLINICAL DERMATOLOGY

Contact allergy

Paraben paradox

The “paraben paradox” is a well-known phenomenon in allergic contact dermatitis.

Parabens are esters of *p*-hydroxybenzoic acid. Methyl, ethyl, propyl, and butyl *p*-hydroxybenzoates are most commonly used in medications, cosmetics, paste bandages, ultrasound gels, dentifrices, suppositories, and foodstuffs. The term “paraben paradox” was first coined by Fisher in 1973.^[2] He described two paradoxical phenomena related to paraben:

- Paraben containing toiletries that are common causes of contact allergy often produce false negative results when patch tested on the back.
- Many individuals who are allergic (patch test positive) to parabens can continue using cosmetics containing them on normal skin without developing any dermatitis, but react to such topical medications when applied over compromised skin (e.g., venous ulcer).

Fisher attributed these phenomena to the insufficiently low concentration of parabens in such cosmetic preparations that fail to produce contact allergy on normal skin and the fact that patch testing is also carried over normal skin. Cashman, *et al.*,^[3] in 2005, proposed the “esterase” and “microbial metabolite” hypotheses to explain the paraben paradox observed by Fisher.

- The esterases present in subcutaneous tissue hydrolyze the long chain parabens (butylparaben) that are least reactive, while the highly reactive short chain parabens (methyl and propylparaben) are metabolized by the esterases in epidermis. A compromised epidermis is therefore associated with altered metabolism and/or increased penetration of such short chain

Access this article online

Website: www.idoj.in

DOI: 10.4103/2229-5178.110589

Quick Response Code:



Address for correspondence:

Dr. Arun C. Inamadar,
Department of
Dermatology,
Venereology and Leprosy,
Shri B. M. Patil Medical
College, Hospital and
Research Center,
Bijapur Liberal
Development Education
University,
Bijapur, Karnataka,
India.
E-mail: aruninamadar@rediffmail.com

highly reactive parabens into the subcutaneous tissue and consequent allergic reaction.

- According to the microbial metabolite hypothesis, repeated application of paraben containing topical preparations leads to selective proliferation of paraben-resistant microbial strains. These organisms hydrolyze parabens to hydroxybenzoic acid, which induce or accentuate sensitization.^[3]

Lanolin paradox

Similar to paraben paradox, the “lanolin paradox” was described by Wolff in 1996.^[4] Lanolin is a natural product obtained from sheep fleece that contains complex mixture of esters and polyesters of high molecular weight alcohols and fatty acids. Lanolin is mostly used in topical medications, emollients, and cosmetics. Lanolin-sensitive patients often are patch test negative to pure lanolin, which may be due to a low concentration of allergens and also the fact that when used “as is,” pure lanolin is a weak sensitizer.^[5,6]

Nail lacquers

Contact allergy to nail lacquers is quite frequent and mostly attributed to the resins included in the products intended to improve the gloss and adhesion of the lacquer film. Tosylamide/formaldehyde resins are most commonly used in lacquers.^[7] Allergic contact dermatitis to these resins may occur on any part that are accessible to the nails (commonly the eyelids, face, neck, and occasionally the thighs and genitalia) and paradoxically in many, no signs of dermatitis are seen at the periungual area.^[8,9]

Nickel Allergy

Among the metals causing contact allergy, nickel is by far the most frequently encountered one, with sensitivity being more common in women than men. The commonest mode of primary nickel sensitization is by prolonged and direct skin contact with corroded nickel-releasing objects. Oral exposure to nickel was also shown to exacerbate the cutaneous dermatitis, and such patients were benefited after reduction of dietary nickel.^[10] However, in some nickel insensitive individuals, oral exposure (e.g., drinking water, dental braces) may induce tolerance.^[11,12] Reduced incidence of cutaneous eruption/exacerbation following low nickel diets in previously sensitized individuals has been reported.^[13] Studies have also shown localized oral as well as cutaneous exacerbation with orthodontic implants in previously sensitized individuals.^[14] On the other hand, lower incidence of oral contact reactions following dental brace implants in nickel-sensitive patients has also been described.^[15] These phenomena of tolerance induction and reduced severity of the disease are attributed to selective induction of a suppressor T-cell mediated immunological tolerance or suppression of hypersensitivity following oral nickel exposure.^[12]

Pigmentary derangements

Vitamin B12 deficiency

Mucocutaneous features of vitamin B12 deficiency include glossitis, angular cheilitis, cutaneous hyperpigmentation, and hair depigmentation. Hyperpigmentation may be diffuse or mottled with predilection for dorsa of hands, palmar creases, knuckles, and flexures. The cause for hyperpigmentation is attributed to decreased levels of reduced glutathione that normally has an inhibitory effect on tyrosinase. Consequently, decreased reduced glutathione is associated with increased tyrosinase activity and melanin synthesis. Histopathology reveals increased number of melanocytes in the basal layer with melanin incontinence into the dermis. Associated hair depigmentation, which may be diffuse or localized, is considered to be a paradox, and Niiyama *et al.*, proposed that the pigmentary incontinence in hair follicles might lead to cavities with resultant depigmentation as opposed to the skin where hyperpigmentation results.^[16,17]

Chronic renal failure

Patients with chronic renal failure (CRF) and/or those on dialysis exhibit a variety of cutaneous pigmentary changes. Pallor of the skin attributed to decreased hematopoiesis and increased hemolysis is the earliest and commonest finding.^[18] Hyperpigmentation, especially over the sun exposed areas, as a result of altered renal metabolism of β -melanocyte-stimulating hormone and yellowish discoloration due to retention of carotenoids and urochromes in the dermis are other cutaneous pigmentary changes associated with CRF and dialysis.^[19] Hypopigmentation of the skin and hair has been described as a paradoxical phenomenon and has been attributed to altered phenylalanine metabolism generating tyrosinase-inhibiting metabolites.^[20-22]

Exogenous ochronosis

Exogenous ochronosis is almost exclusively described in blacks, especially in African women. Although hydroquinone containing medications are widely used in the USA, the condition is very uncommon there. This paradoxical sparing of white skin is possibly due to the absence of factors seen in blacks such as a higher enzyme activity associated with melanin synthesis and intense ultraviolet (UV) exposure.^[23]

Leucoderma in melanoma

Vitiligo-like leucoderma in melanoma is a well known, yet infrequent phenomenon and is especially seen in patients on immunotherapy.^[24] Although vitiligo-like lesions are associated with an advanced stage of the disease with metastases, it is paradoxically associated with good prognosis.^[25] Development of leucoderma in melanoma may be limited to the lesion, appear as a halo around the lesion, or be widespread and possibly indicates a T-cell mediated immunological response^[26] that targets antigens present both on melanoma cells and on

normal melanocytes as well (e.g., MART-1, tyrosinase, gp100, tyrosinase-related protein 1 and 2),^[27] thus resulting in melanoma regression and vitiligo.

Others

Hypothyroidism

Hair changes in acquired hypothyroidism are predominantly characterized by diffuse thinning and sparseness of body and scalp hair. Loss of lateral one-third of the eyebrows (Hertog's sign) is a characteristic finding as well. In some patients, especially among children, a paradoxical hypertrichosis characterized by growth of coarse terminal hair is seen over the shoulders, back, and outer surface of the lower limbs.^[28,29] Although the exact mechanism is unclear, several hypotheses like associated hyperprolactinemia,^[30] prolongation of the anagen,^[31] and an associated keratinization abnormality^[32] have been proposed.

Lichen planus

Itching is a consistent symptom of Lichen planus (LP) and may range from mild to obdurate. Itching is particularly present during the early eruptive stage and in hypertrophic LP. Paradoxically, excoriations are conspicuously absent as the patients rubs rather than scratches to gain relief and, hence, the pathognomonic lesional morphology and Wickham's striae all are well preserved.^[33,34]

Bleeding diathesis in myeloproliferative disorders

The myeloproliferative disorders^[35] including essential thrombocythemia, polycythemia vera, and primary myelofibrosis are associated with an abnormally increased platelet count that clinically manifests commonly with thrombotic episodes affecting various organs. Paradoxically, a bleeding diathesis may also be encountered, especially with higher platelet counts ($>1000 \times 10^9/L$)^[36] that may range from ecchymoses or mucosal bleeds to dreadful internal hemorrhage. This is due to an acquired von Willebrand disease (loss of large von Willebrand factor multimers that results in its functional defect) and/or qualitative platelet defects (acquired storage pool disorder).^[37]

Familial occurrence of sporadic diseases

Certain genodermatoses that almost exclusively occur sporadically have been unusually described to be occurring in a familial pattern affecting different generations. Several explanations like multifactorial etiology^[38] or an autosomal dominant inheritance with variable expressivity^[39] have been proposed for such paradoxical occurrence. In 1993, Happle described the concept of paradominant inheritance pattern to explain such observations.^[40] Accordingly, individuals heterozygous for a "paradominant trait" transmit the mutated gene through several generations imperceptibly, and the

condition manifests clinically only when an additional somatic mutation occurs in the early embryogenesis, which leads to loss of heterozygosity. Few examples for such paradominant traits include Becker's nevus syndrome,^[41] Klippel-Trenaunay syndrome,^[40,42] large congenital melanocytic nevi,^[43] congenital triangular alopecia,^[44] and some vascular malformations.

Melanoma and sun exposure

Sun exposure is undoubtedly a significant risk factor for the development of cutaneous melanoma. However, the incidence or risk of development of melanoma is determined by the type of sun exposure. Intermittent, recreational exposure, and a history of sunburns in childhood are perhaps more important in melanoma causation rather than continuous, occupational exposure without history of sunburn. Paradoxically, continuous exposure has been shown to be beneficial or be protective against melanoma development. Photoadaptation (increased melanization and epidermal thickening) and protective effects of vitamin D are possibly associated with continuous solar exposure that reduces melanoma risk.^[45,46] The mortality in melanomas associated with chronic sun exposure is also paradoxically less as compared to those associated with no such exposure.^[47]

Neuroblastoma

Although the stage IV-S neuroblastoma^[48] is associated with distant metastases, the prognosis, however, is paradoxically favorable. The tumor may remit spontaneously^[49] or transform into a benign ganglioneuroma.^[50] Based on this biological behavior of the tumor, infants with stage IV-S disease are best observed carefully with minimal intervention. The mortality in these patients has been attributed to the complications secondary to the massive hepatomegaly leading to respiratory distress, and disseminated intravascular coagulation. It is only under these circumstances that an aggressive treatment approach is warranted.^[51] Studies have also shown that complete resection of the primary tumor does not alter the overall outcome of the disease.^[52]

IMMUNOLOGICAL PARADOXES

Human immunodeficiency virus (HIV) infection and AIDS

Target cell and viral load

The CD4+ T-lymphocytes are the target cells for the HIV. A paradoxical feature of long-term HIV-1 infection is the inverse relationship between the levels of these CD4 T-cells and the viral load in the blood.^[53] With advancing depletion in the target CD4+ cells, there is increase in the plasma viral load. This paradox could be explained by increased viral production per infected CD4+ cell in the setting of a waning (possibly CD8+ mediated) immune surveillance.^[54] Also, the tissue macrophages serve as an alternate reservoir in which the HIV replicate even with declining CD4+ cell count.^[55]

Hyperimmune conditions in AIDS

Various “AIDS defining” illnesses like oral hairy leukoplakia, giant molluscum contagiosum, bacillary angiomatosis, systemic mycoses, and atypical mycobacterial infections occur when the CD4+ T-cell count falls below 200/ μ L and are indicative of profound immunodeficiency. However, at this stage, several non-infectious inflammatory hyperimmune dermatoses including eosinophilic folliculitis, intense hypersensitivity to insect bites (manifesting as pruritic papular eruptions), seborrheic dermatitis, and drug hypersensitivity are also encountered.^[56] The exact cause of this paradox has not been delineated, but a T-helper-2 (T_H2) immune response that predominates in progressive HIV disease might possibly facilitate the development of such disorders.^[57,58]

HIV and human papilloma virus

Patients seropositive for HIV have more extensive and high-risk human papilloma virus (HPV) associated diseases. This is attributed to the progressive decline in the CD4+ T-cells that allow an increased prevalence of HPV-related infections, type-specific persistence, and an increase in HPV-associated malignancies. Although highly active anti-retroviral therapy (HAART) has a positive impact on patients' lives by reconstituting CD4+ T-cell count and decreasing the viral load, paradoxically however, the incidence of HPV-induced diseases has increased rather than decreased since the introduction of HAART.^[59-62] This may partly be due to increased exposure to diverse HPV types, particularly high-risk types that persist longer in anogenital regions, to this upgrading immunity. Also, persistent or emergent HPV disease in HIV infection might represent persistent or modulated immune dysregulation after HAART initiation and, hence, may be viewed as a form of immune reconstitution associated disease.^[63]

Psoriasis and HIV

Psoriasis, a T-cell mediated dermatosis that responds to T-cell immunosuppressants (e.g., cyclosporine), is seen to exacerbate or occur in severe or recalcitrant forms in HIV infected patients.^[64] Furthermore, with advancing immunosuppression and declining CD4+ T-cell counts, psoriasis and psoriatic arthropathy worsen and improve with HAART initiation.^[65] This paradoxical exacerbation is possibly due to loss of regulatory T-cells and increased activity of CD8+ T-cells.^[66] Opportunistic infections, especially candidiasis and staphylococcal infections, or drugs may also contribute to the flare of psoriasis in HIV infected patients.^[64]

Leprosy and HIV

Leprosy in HIV infected patients behaves no differently than in the non-infected individuals. In contrast to tuberculosis, the natural course leprosy is not altered with HIV co-infection. The clinical, immunological, and pathological features are the same even in progressive HIV disease and the granuloma formation,

in contrast to tuberculosis, remains well-preserved even with declining T-cell mediated immunity—the so-called “granuloma paradox”. This paradox may be due to the differences between the activation state and rates of cell turnover within leprosy and tuberculosis granulomas that differentially affect the susceptibility of the granulomas to HIV.^[67-69]

Others

Leprosy and host T-cell immunity

After acquiring the infection, the pathogenesis of leprosy and the consequent clinical manifestations are largely determined by the extent of host T-cell mediated immunity mounted against *Mycobacterium leprae*. In children, the indeterminate, tuberculoid, and borderline tuberculoid forms predominate and, paradoxically, multibacillary disease is rare even in those with poor cell-mediated immunity.^[70]

Phenomenon of compartmentalization

Compartmentalization of inflammatory cells in the peripheral tissues is one of the proposed mechanisms for peripheral anergy seen in some immune-mediated disorders. In the following instances, although an intense immunological activity occurs in the involved tissues, there is however, a global reduction in immune response to antigens (anergy) or peripheral immune-cytopenia.

1. Sarcoidosis

Sarcoidosis is a multisystem disorder in which involved organs and tissues exhibit a common pathological change, the non-caseating granuloma, mediated by CD4 + helper T-cells of the T_H1 subtype. On the one hand, affected organs express intense immune response, while, on the other, there is a globally decreased delayed type hypersensitivity to common antigens (anergy). This paradox is particularly evident during the active phase of the disease. Compartmentalization of monocytes and T-lymphocytes in the peripheral tissues is one of the proposed mechanisms for peripheral lymphopenia and anergy.^[71] A recent study showed that, in sarcoidosis, there is an expansion of regulatory T-cells (CD4+, CD25^{bright}) that suppress naive T-cell proliferation, leading to anergy, but are unable to completely suppress tumor necrosis factor alpha (TNF- α), which is principally involved in the disease pathogenesis.^[72]

2. Psoriasis

In similarity with the phenomenon of compartmentalization in sarcoidosis, psoriasis is characterized by reduced levels of circulating natural killer (NK) cells^[73] as they are dedicated to in the pathogenesis of the disease, which is evidenced by increased levels of *perforin* (cytolytic product of the NK cells) in the epidermis of psoriatic plaques.^[74] A similar phenomenon is also seen in other autoimmune diseases like rheumatoid arthritis.^[73]

Androgens and hair growth

Hair growth in human beings is influenced by the circulating androgen levels. However, this influence is not uniform for all the hairs. At puberty, the beard and other secondary sexual hairs grow, the scalp hair is gradually lost, while the eyebrows and eyelashes remain unchanged.^[75] This paradoxical effect of androgen at different hair bearing sites is attributed to the differential response of the follicular dermal papillae to androgen stimulation in these sites.^[76] In response to androgen stimulation, the dermal papillae in the beard area secrete various autocrine growth factors, including the insulin-like growth factor 1 (IGF1), which has a stimulatory effect on the follicles.^[77] On the scalp, however, these papillae secrete transformation growth factor beta1 (TGF- β 1) that has an inhibitory effect on the hair follicles. This TGF- β 1 mediated activity is maximum in the anterior aspect of the scalp, especially so, over the temples and hence, the development of "patterned" hair loss.^[78]

Immune dysregulation in ataxia telangiectasia

Ataxia telangiectasia (AT) is a rare autosomal recessive disorder due to mutations in *ATM* (ataxia telangiectasia mutated) gene that encodes a phosphatidyl kinase involved in cell cycle control and DNA repair. AT is characterized by progressive neurodegeneration, oculocutaneous telangiectasia, variable immunodeficiency, and a high predisposition for lymphoreticular malignancies. Most common immunodeficiency profile, especially in those with complete absence of *ATM* gene activity, is that of IgA deficiency (especially IgA2) and lymphopenia (both T and B cells, except NK cells). Paradoxically, IgM, IgA, and IgG can be elevated in some patients, including monoclonal gammopathy in more than 10% of cases. The *ATM* gene also plays a role in immunoglobulin class switch recombination (Ig-CSR), which may be the cause for elevation of IgM. Concurrent infections may be also be related to such elevations.^[79,80]

Psoriasis and pregnancy

Being a T-cell (T_H1) mediated disorder, psoriasis in general, has been shown to improve during pregnancy, which is associated with a shift of T_H1 to T_H2 cytokine profile.^[81] This shift is attributed to elevated estrogen^[82] and progesterone^[81,83] individually (as both are known to have immunosuppressive functions) and to the altered estrogen progesterone ratio^[84] in general as well. Paradoxically, some pregnancies are associated either with no change or even worsening of the disease, which has been postulated to be due to the achievement of lower levels of elevation in estrogen relative to progesterone as compared to those pregnancies in which psoriasis improves.^[82] Also, impetigo herpetiformis, a type of pustular psoriasis is induced by pregnancy.

Lupus anticoagulant

Antiphospholipid syndrome (APS) is defined as per the

international consensus statement preliminary criteria for antiphospholipid antibody syndrome.^[36,85] The syndrome is mainly characterized by thrombotic events, recurrent pregnancy loss, and thrombocytopenia that are associated with antiphospholipid antibodies in the serum including anti-cardiolipin, anti- β 2-glycoprotein I (β_2 GPI) antibodies, and lupus anticoagulant (LA). Several mechanisms have been proposed for the development of thrombosis such as promotion of coagulation reactions, interference with anticoagulant or fibrinolytic pathways, and platelet activation.^[36] The current consensus on the mechanism leading to the development of thrombosis in APS is the one mediated by anti- β_2 GPI antibodies. Binding of anti- β_2 GPI antibodies leads to the disruption of a crystal shield of *annexin V* that covers the platelet membrane and normally prevents binding of procoagulant molecules to its surface.^[36,86] The LA however is associated with thrombosis *in vivo*, whereas, *in vitro*, is seen to prolong the coagulation tests that are employed to detect it and hence its name. The exact mechanism of this paradox as to how an "anticoagulant" promotes thrombosis *in vivo* remains to be clearly elucidated.^[87]

PARADOXES IN DERMATOPATHOLOGY

Clinical-pathological disparities

Pauci-inflammatory photodermatitis

Polymorphic light eruption (PLE) is most common among the idiopathic photodermatoses. It is common in the first decades of life with females being more affected than males. As the name suggests, PLE manifests in different morphological patterns (papular, papulovesicular, erythematous, eczematous, or plaque like).^[88] Prominent histopathological features include epidermal spongiosis and exocytosis with perivascular lymphoid infiltrate that is milder in early lesions and moderate-to-intense in late lesions. Mural and endothelial edema is also noted in the dermal vessels.^[89] In some cases, clinically obvious lesions are discordantly associated with no or minimal histopathological changes. Such paradoxical photosensitive lesions are described under pauci-inflammatory photodermatitis.^[90]

Sebaceous glands in aging skin

In the elderly, in spite of decreased output from sebaceous glands, which is attributed to decreased adrenal and gonadal androgen synthesis,^[91] their size however, is increased owing to decreased cell turnover.

Clinical course/picture versus histopathology

Certain conditions exhibit considerable discrepancy between their clinical behavior and histological features. A seemingly benign histological profile of a condition may be associated with a paradoxically aggressive clinical behavior or significant morbidity and vice-versa. Following are examples of such conditions:

Infantile fibromatosis

Infantile fibromatosis is one of the locally recurring fibromatoses that is histologically characterized by homogenous proliferation of bundles of spindle shaped cells with bland nuclei, indicating entirely banal nature of the lesion. The clinical behavior is however that of a rapidly growing tumor with potential to reach significant size and entrap vascular, neural, and articular structures, leading to morbidity. They are also persistent and promptly recur following excision. However, unlike fibrosarcoma, the lesions do not metastasize. Similarly, nodular fasciitis, a reactive myofibroblastic proliferation shows completely benign histopathological features, but paradoxically, it is a more rapidly growing lesion than any other potentially more aggressive process.^[92]

Myofibromatosis

Myofibromatosis is characterized by multiple lesions affecting the subcutaneous tissue as well as the viscera that may be associated with a poor prognosis, not because of its clinical and pathological behavior (which is totally benign), but due to its sheer numbers in vital organs that compromise their function with possible fatality.^[92]

Lymphomatoid papulosis

Lymphomatoid papulosis (LyP) is a condition characterized by a benign clinical picture, but, histopathologically shows features of cutaneous (CD30+) lymphoma. LyP is characterized by papules papulonecrotic or papulonodular lesions, affecting predominantly the upper trunk and proximal limbs of adult females, which are otherwise innocuous. The characteristic feature of LyP is the occurrence of the lesions in recurrent crops that typically evolve rapidly within a few days and resolve spontaneously over few weeks to months, leaving behind atrophic scarring in many. Such crops recur every few months without any obvious inciting factors. Histopathologically, LyP is characterized by the presence of atypical lymphocytes with large nuclei and abnormal mitoses. Based on the predominant population of the cells, LyP is grouped into type A (Hodgkin's lymphoma like), type B (mycosis fungoides like), and type C (large anaplastic B cell lymphoma like).^[93] Approximately, 5-10% of LyP progress to malignant lymphoma, mostly to mycosis fungoides.

Others

Basal cell carcinoma

Basal cell carcinoma (BCC) is the most common neoplasm occurring in humans. It is a neoplasm arising from the basal cells of the epidermis and also from outer root sheath of pilosebaceous units in a minority. Histopathologically, though numerous mitoses^[94] are present, clinically the tumor is very slow growing. This paradox has been attributed to the correspondingly high apoptotic index with low expression of *survivin*, an apoptosis inhibitor in BCCs.^[95]

Aging skin

In chronologically aging skin, the elastic fibers gradually lessen, most likely due to enzymatic degradation of elastin.^[91] Together with decreased elastin, the skin becomes wrinkled, saggy, and inelastic, due to decreased collagen synthesis. Paradoxically, the photo-aged skin shows focally increased elastin owing to UV-induced transcriptional activation of the elastin gene^[96] and lysozyme-induced reduction in leukocyte elastase activity.^[97] Together with heliodermatitis,^[91] photo-aged skin appears thick, dry, rough, deeply wrinkled, and inelastic owing to the disarrangement of collagen and elastic fibers.^[98]

PARADOXES IN DERMATOLOGICAL THERAPY

Paradoxical induction/exacerbation of dermatoses

Antihistamine-induced urticaria

Antihistamines are the first-line treatment for urticaria, which are generally quite safe, allowing long-term administration. Paradoxically, antihistamine-induced urticaria, which, although uncommon, has been encountered, leading to confusion in the clinical setting. Lee, *et al.*, reviewed 16 reports on antihistamine-induced urticaria that showed cetirizine being the most commonly reported culprit. The exact mechanism of this paradox is not clear. Type I and IV hypersensitivity reactions to antihistamines, non-immune mast cell activation-degranulation, and intolerance to antihistamines have been speculated.^[99]

Anti-TNF- α biological induced psoriasis

The anti-TNF- α biologicals are being used to treat many inflammatory and autoimmune dermatoses. They (infliximab and etanercept) are especially used for recalcitrant extensive plaque psoriasis, generalized pustular psoriasis, and psoriatic arthritis. Cutaneous adverse effects of these agents include lichenoid, eczematous, urticarial, and other forms of lesions.^[100] One of the perplexing side effects of these agents is the induction and/or exacerbation of psoriasis itself. There are many published reports of such anti-TNF biological induced (new-onset psoriasis in absence of past or family history) as well as exacerbated psoriasis (plaque and pustular types).^[100-103] The exact pathophysiology of this paradox is unclear and, moreover, such psoriatic lesions are identical both clinically and histologically to idiopathic psoriasis. Unhindered interferon- α (IFN- α) release by plasmacytoid dendritic cells (PDC) in the background of TNF- α inhibition is one of the proposed mechanisms. Increased IFN- α also induces expression of CXCR-3 chemokine receptor on T-cells that facilitates their homing to the skin, which is also considered to be a mechanism of psoriasis induction or exacerbation in patients receiving anti-TNF- α biologicals.^[104,105]

Anti-TNF- α biological-induced vasculitis

Infliximab has been used in the treatment of refractory

systemic vasculitis with great benefits.^[106] However, anti-TNF- α biologicals have been paradoxically reported to induce vasculitis in patients with rheumatic diseases. In a recent study, cutaneous as well as systemic (renal vasculitis and peripheral neuropathy) vasculitis developed in equal number of patients treated with anti-TNF- α agents for rheumatoid arthritis and inflammatory bowel disease. As these diseases themselves may be associated with cutaneous vasculitis, the authors laid down criteria so as to attribute the development of vasculitis to the anti-TNF- α agents administered.^[107] Although the exact cause of such vasculitis is unclear, development of auto-antibodies against these anti-TNF- α agents with subsequent immune complex deposition and inflammation is one of the proposed mechanisms.^[108,109] Another hypothesis is the increased IFN- α (see above) mediated induction of autoimmune diseases like lupus erythematosus dermatomyositis including vasculitis.^[110]

Anti-TNF- α agent-induced sarcoidosis

Similar to vasculitis, the anti-TNF- α agents are useful in refractory sarcoidosis and here as well, these agents may paradoxically induce sarcoid-like granulomas (cutaneous and pulmonary). The mechanism of such granuloma development has again been attributed to increased IFN- α , which mediates granuloma formation to the antigenic stimuli, possibly *M. tuberculosis* (pulmonary) and *P. acnes* (cutaneous).^[111]

Isotretinoin and acne fulminans

Acne fulminans is a severe form of acne with systemic features, that may occur in some predisposed individuals as a hypersensitivity reaction to *P. acnes*.^[112] The treatment of choice is a sequential therapy beginning with systemic corticosteroids to combat the early crisis followed by addition of low-dose oral isotretinoin for 3-4 months with the steroids being tapered off gradually. However, isotretinoin must be used with caution, as paradoxical induction/exacerbation of acne fulminans has been reported.^[113-115]

Topical calcineurin inhibitors (TCIs)-induced rosacea

The TCIs like tacrolimus and pimecrolimus are relatively newer topical treatment modalities used for the treatment of rosacea^[116] and have been found to be both effective and safe. However, reports of rosaceiform dermatitis following long-term usage of these agents have also been documented. Topical immunosuppressive effects of the TCIs leading to overgrowth of *Demodex* mites in skin together with their inherent vasoactive properties possibly act synergistically leading to "iatrogenic rosacea."^[117-119]

D-penicillamine-induced pseudoscleroderma

D-penicillamine is used as an antifibrotic agent in treatment of systemic sclerosis, as it is a known lathyrogen that acts by interfering with cross-linking of collagen. It is generally recommended in early disease with progressive skin changes,

pulmonary compromise, or renal disease.^[120] Paradoxically, morphea-like cutaneous lesions (pseudoscleroderma) have been reported in patients receiving D-penicillamine for other indications.^[121,122]

Paradoxical antagonistic effects

H1-antihistamine-induced seizures

The H1-antihistamines are widely used for various pruritic conditions. These are traditionally grouped into first generation and second generation H1-antihistamines, based on their ability to produce sedation. The first generation H1-antihistamines cross the blood-brain barrier and hence, are sedative. In infants and children, these "sedative" first generation H1-antihistamines may paradoxically precipitate seizures, especially in known epileptics or cases of febrile convulsions.^[123] In infants and children, the histaminergic system is probably a more important inhibitory system in the central nervous system than the γ -aminobutyrate (GABA-nergic) system with regard to convulsions. Hence, the first generation H1-antihistamines may reduce the threshold for seizure precipitation by blocking the histamine activity.^[124]

Q-switched laser-induced tattoo darkening

The Q-switched lasers are currently the lasers of choice for the treatment of tattoos. Although post-treatment hyperpigmentation is not infrequently encountered, it is generally transient. Paradoxical darkening of the tattoos, especially the red, white, or flesh-toned ones occur occasionally and the patients must be informed beforehand about this. These tattoo pigments often contain ferric oxide (rust-brown color) and titanium dioxide (white color). Following laser treatment, these pigments get reduced to ferrous oxide (black color) and blue-colored titanium dioxide respectively, resulting in immediate post-treatment darkening of such tattoos.^[125]

Intense pulsed light (IPL)-induced hypertrichosis

Unwanted hair removal with lasers and IPL is becoming widely popular as it is both effective and safe. Choosing an appropriate treatment modality (laser or IPL), appropriate wavelength, and appropriate pulse width and fluence to deliver the wavelength at, are all imperative for optimal treatment outcome, as paradoxical hypertrichosis has been observed after laser treatment, especially in dark-skinned individuals. This occurrence has been attributed to the suboptimal fluences that possibly stimulate the follicles.^[126]

REFERENCES

1. Oxford dictionaries. Available from: <http://oxforddictionaries.com/definition/english/paradox> [Last accessed on 2013 Jan 20].
2. Fisher AA. The paraben paradox. *Cutis* 1973;12:830-2.
3. Cashman AL, Warshaw EM. Parabens: A review of epidemiology, structure, allergenicity, and hormonal properties. *Dermatitis* 2005;16:57-66.
4. Wolf R. The lanolin paradox. *Dermatology* 1996;192:198-202.

5. Lee B, Warshaw E. Lanolin allergy: History, epidemiology, responsible allergens, and management. *Dermatitis* 2008;19:63-72.
6. Cronin E. Lanolin dermatitis. *Br J Dermatol* 1966;78:167-74.
7. De Berkar DA, Baran R. Disorders of nails. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 65.1-65.57.
8. Lidén C, Berg M, Färm G, Wrangsjö K. Nail varnish allergy with far-reaching consequences. *Br J Dermatol* 1993;128:57-62.
9. White JM, de Groot AC, White IR. Cosmetics and skin care products. In: Johansen JD, Frosch PJ, Lepoittevin JP, editors. *Contact Dermatitis*. 5th ed. Berlin: Springer; 2011. p. 591-605.
10. Veien NK, Hattel T, Laurberg G. Low nickel diet: An open, prospective trial. *J Am Acad Dermatol* 1993;29:1002-7.
11. White JM, Goon AT, Jowsey IR, Basketter DA, Mak RK, Kimber I, et al. Oral tolerance to contact allergens: A common occurrence? A review. *Contact Dermatitis* 2007;56:247-54.
12. Van IM, Andersen KE, Von Blomberg BM, Boden D, Bruynzeel DP, Burrows D, et al. Reduced frequency of nickel allergy upon oral nickel contact at an early age. *Clin Exp Immunol* 1991;85:441-5.
13. Santucci B, Cristaudo A, Cannistraci C, Picardo M. Nickel sensitivity: Effects of prolonged oral intake of the element. *Contact Dermatitis* 1988;19:202-5.
14. Schultz JC, Connelly E, Glesne L, Warshaw EM. Cutaneous and oral eruption from oral exposure to nickel in dental braces. *Dermatitis* 2004;15:154-7.
15. Kerosuo H, Kullaa A, Kerosuo E, Kanerva L, Hensten-Pettersen A. Nickel allergy in adolescents in relation to orthodontic treatment and piercing of ears. *Am J Orthod Dentofacial Orthop* 1996;109:148-54.
16. Jen M, Shah KN, Yan AC. Cutaneous changes in nutritional diseases. In: Wolff K, Goldsmith LA, Katz SI, Gilchrist BA, Paller AS, Leffell DJ, editors. *Fitzpatrick's Dermatology in General Medicine*. 7th ed. New York: The McGraw-Hill Companies; 2008. p. 1201-18.
17. Niiyama S, Mukai H. Reversible cutaneous hyperpigmentation and nails with white hair due to vitamin B12 deficiency. *Eur J Dermatol* 2007;17:551-2.
18. Cox NH, Coulson IH. Systemic disease and the skin. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 62.1-62.113.
19. Udayakumar P, Balasubramanian S, Ramalingam KS, Lakshmi C, Srinivas CR, Mathew AC. Cutaneous manifestations in patients with chronic renal failure on hemodialysis. *Indian J Dermatol Venereol Leprol* 2006;72:119-25.
20. Gharibi L, Zouhair K, Ramdani B, Benchikhi H. Hair lightening in an hemodialysis patient treated by heptaminol (heptamyl). *Dermatol Online J* 2009;15:16.
21. Lapeere H, Boone B, de Schepper S, Verhaeghe E, Ongenae K, Van Geel N, et al. Hypomelanoses and hypermelanoses. In: Wolff K, Goldsmith LA, Katz SI, Gilchrist BA, Paller AS, Leffell DJ, editors. *Fitzpatrick's Dermatology in General Medicine*. 7th ed. New York: The McGraw-Hill Companies; 2008. p. 622-40.
22. Hmida MB, Turki H, Hachicha J, Reygagne P, Rabier D, Zahaf A, et al. Hypopigmentation in hemodialysis. Acquired hair and skin fairness in a uremic patient undergoing maintenance hemodialysis: Case report and review of the literature. *Dermatology* 1996;192:148-52.
23. Berth-Jones J. Topical therapy. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 73.1-73.52.
24. Daneshpazhooh M, Shokoohi A, Dadban A, Raafat J. The course of melanoma-associated vitiligo: Report of a case. *Melanoma Res* 2006;16:371-3.
25. Wang JR, Yu KJ, Juan WH, Yang CH. Metastatic malignant melanoma associated with vitiligo-like depigmentation. *Clin Exp Dermatol* 2009;34:209-11.
26. Becker JC, Guldberg P, Zeuthen J, Bröcker EB, Straten PT. Accumulation of identical T cells in melanoma and vitiligo-like leukoderma. *J Invest Dermatol* 1999;113:1033-8.
27. Cunha D, Pacheco FA, Cardoso J. Vitiligo: A good prognostic factor in melanoma? *Dermatol Online J* 2009;15:15.
28. Perloff WH. Hirsutism-a manifestation of juvenile hypothyroidism. *JAMA* 1955;157:651-2.
29. Stern SR, Kelnar CJ. Hypertrichosis due to primary hypothyroidism. *Arch Dis Child* 1985;60:763-6.
30. Sharma D, Shanker V, Tegta G, Gupta M, Verma GK. Clinico-investigative profile of patients of hirsutism in a tertiary level institution. *Int J Trichology* 2012;4:69-74.
31. Camcho-Martínez FM. Hypertrichosis. In: Blume-Peytavi U, Tosti A, Whiting DA, Trübe R, editors. *Hair Growth and Disorders*. 1st ed. Berlin: Springer; 2008. p. 333-56.
32. Maekawa Y. Rolled hair and hypertrichosis: A manifestation of juvenile hypothyroidism. *J Dermatol* 1983;10:157-60.
33. Berathnach SM. Lichen planus and lichenoid disorders. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 41.1-41.28.
34. Reich A, Welz-Kubiak K, Szepletowski JC. Pruritus differences between psoriasis and lichen planus. *Acta Derm Venereol* 2011;91:605-6.
35. Campbell PJ, Green AR. The myeloproliferative disorders. *N Engl J Med* 2006;355:2452-66.
36. Cox NH, Piette WW. Purpura and microvascular occlusion. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 49.1-49.51.
37. Deshmukh A, Subbiah SP, Malhotra S, Deshmukh P, Pasupuleti S, Mohiuddin S. Spontaneous hemopericardium leading to cardiac tamponade in a patient with essential thrombocythemia. *Cardiol Res Pract* 2011;2011:247814.
38. Aelvoet GE, Jorens PG, Roelen LM. Genetic aspects of the Klippel-Trenaunay syndrome. *Br J Dermatol* 1992;126:603-7.
39. Blei F, Walter J, Orlow SJ, Marchuk DA. Familial segregation of hemangiomas and vascular malformations as an autosomal dominant trait. *Arch Dermatol* 1998;134:718-22.
40. Happle R. Klippel-Trenaunay syndrome: Is it a paradominant trait? *Br J Dermatol* 1993;128:265.
41. Luna PC, Nocito MJ, Cañadas NG, Posse ML, Marchesi C, Lustia MM, et al. Becker nevus syndrome. Report of three cases. *Dermatol Argent* 2008;14:379-82.
42. Hofer T, Frank J, Itin PH. Klippel-Trenaunay syndrome in a monozygotic male twin: Supportive evidence for the concept of paradominant inheritance. *Eur J Dermatol* 2005;15:341-3.
43. Danarti R, König A, Happle R. Large congenital melanocytic nevi may reflect paradominant inheritance implying allelic loss. *Eur J Dermatol* 2003;13:430-2.
44. Happle R. Congenital triangular alopecia may be categorized as a paradominant trait. *Eur J Dermatol* 2003;13:346-7.
45. Newton-Bishop JA, Chang YM, Elliott F, Chan M, Leake S, Karpavicius B, et al. Relationship between sun exposure and melanoma risk for tumours in different body sites in a large case-control study in a temperate climate. *Eur J Cancer* 2011;47:732-41.
46. Parkin DM, Mesher D, Sasieni P. Cancers attributable to solar (ultraviolet) radiation exposure in the UK in 2010. *Br J Cancer* 2011;105 (Suppl 2):S66-9.
47. Corona R. Another paradigm lost or just a paradox? *Arch Dermatol* 2005;141:1587-8.
48. Neuroblastoma. Available from: <http://en.wikipedia.org/wiki/Neuroblastoma> [Last accessed on 2013 Feb 1].
49. Rangelcroft L, Lauder I, Wagget J. Spontaneous maturation of stage IV-S neuroblastoma. *Arch Dis Child* 1978;53:815-7.
50. Evans AE, Baum E, Chard R. Do infants with stage IV-S neuroblastoma need treatment? *Arch Dis Child* 1981;56:271-4.
51. Nickerson HJ, Matthay KK, Seeger RC, Brodeur GM, Shimada H,

- Perez C, et al. Favorable biology and outcome of stage IV-S neuroblastoma with supportive care or minimal therapy: A Children's Cancer Group study. *J Clin Oncol* 2000;18:477-86.
52. Guglielmi M, De Bernardi B, Rizzo A, Federici S, Boglino C, Siracusa F, et al. Resection of primary tumor at diagnosis in stage IV-S neuroblastoma: Does it affect the clinical course? *J Clin Oncol* 1996;14:1537-44.
 53. Harrington PR, Swanstrom R. The biology of HIV, SIV, and other lentiviruses. In: Holmes KK, Sparling PF, Stamm WE, Piot P, Wasserheit JN, Corey L, et al. editors. *Sexually Transmitted Diseases*. 4th ed. New York: The McGraw-Hill Companies; 2008. p. 323-39.
 54. Jin X, Bauer DE, Tuttleton SE, Lewin S, Gettie A, Blanchard J, et al. Dramatic rise in plasma viremia after CD8(+) T cell depletion in simian immunodeficiency virus-infected macaques. *J Exp Med* 1999;189:991-8.
 55. Igarashi T, Brown CR, Endo Y, Buckler-White A, Plishka R, Bischofberger N, et al. Macrophage are the principal reservoir and sustain high virus loads in rhesus macaques after the depletion of CD4+T cells by a highly pathogenic simian immunodeficiency virus/HIV type 1 chimera (SHIV): Implications for HIV-1 infections of humans. *Proc Natl Acad Sci U S A* 2001;98:658-63.
 56. Colven R. Generalized cutaneous manifestations of STD and HIV infection: Typical presentations, differential diagnosis and management. In: Holmes KK, Sparling PF, Stamm WE, Piot P, Wasserheit JN, Corey L, et al. editors. *Sexually Transmitted Diseases*. 4th ed. New York: The McGraw-Hill Companies; 2008. p. 1177-97.
 57. Saavedra A, Johnson RA. Cutaneous manifestations of human immunodeficiency virus disease. In: Wolff K, Goldsmith LA, Katz SI, Gilchrist BA, Paller AS, Leffell DJ, editors. *Fitzpatrick's Dermatology in General Medicine*. 7th ed. New York: The McGraw-Hill Companies; 2008. p. 1927-40.
 58. Amerio P, Verdolini R, Proietto G, Feliciani C, Toto P, Shivji G, et al. Role of Th2 cytokines, RANTES and eotaxin in AIDS-associated eosinophilic folliculitis. *Acta Derm Venereol* 2001;81:92-5.
 59. Van der Burg SH, Palefsky JM. Human Immunodeficiency Virus and Human Papilloma Virus-why HPV-induced lesions do not spontaneously resolve and why therapeutic vaccination can be successful. *J Transl Med* 2009;7:108.
 60. De Vuyst H, Lillo F, Broutet N, Smith JS. HIV, human papillomavirus, and cervical neoplasia and cancer in the era of highly active antiretroviral therapy. *Eur J Cancer Prev* 2008;17:545-54.
 61. Handisurya A, Rieger A, Bankier A, Koller A, Salat A, Stingl G, et al. Human papillomavirus type 26 infection causing multiple invasive squamous cell carcinomas of the fingernails in an AIDS patient under highly active antiretroviral therapy. *Br J Dermatol* 2007;157:788-94.
 62. Fife KH, Wu JW, Squires KE, Watts DH, Andersen JW, Brown DR. Prevalence and persistence of cervical human papillomavirus infection in HIV-positive women initiating highly active antiretroviral therapy. *J Acquir Immune Defic Syndr* 2009;51:274-82.
 63. Meys R, Gotch FM, Bunker CB. Human papillomavirus in the era of highly active antiretroviral therapy for human immunodeficiency virus: An immune reconstitution-associated disease? *Br J Dermatol* 2010;162:6-11.
 64. Duvic M, Johnson TM, Rapini RP, Freese T, Brewton G, Rios A. Acquired immunodeficiency syndrome-associated psoriasis and Reiter's syndrome. *Arch Dermatol* 1987;123:1622-32.
 65. Obuch ML, Maurer TA, Becker B, Berger TG. Psoriasis and human immunodeficiency virus infection. *J Am Acad Dermatol* 1992;27:667-73.
 66. Gudjonsson JE, Johnston A, Sigmundsdottir H, Valdimarsson H. Immunopathogenic mechanisms in psoriasis. *Clin Exp Immunol* 2004;135:1-8.
 67. Kwobah CM, Wools-Kaloustian KK, Gitau JN, Siika AM. Human immunodeficiency virus and leprosy coinfection: Challenges in resource-limited setups. *Case Rep Med* 2012;2012:698513.
 68. Pereira GA, Stefani MM, Araújo Filho JA, Souza LC, Stefani GP, Martelli CM. Human immunodeficiency virus type 1 (HIV-1) and *Mycobacterium leprae* co-infection: HIV-1 subtypes and clinical, immunologic, and histopathologic profiles in a Brazilian cohort. *Am J Trop Med Hyg* 2004;71:679-84.
 69. Ustianowski AP, Lawn SD, Lockwood DN. Interactions between HIV infection and leprosy: A paradox. *Lancet Infect Dis* 2006;6:350-60.
 70. Dogra S, Kanwar AJ. Leprosy (Hansen disease). In: Irvine A, Hoeger P, Yan A, editors. *Harper's Textbook of Pediatric Dermatology*. 3rd ed. Oxford: Wiley-Blackwell; 2011. p. 70.1-70.14.
 71. Howard A, White CR Jr. Non-infectious granulomas. In: Bologna JL, Jorizzo JL, Rapini RP, editors. *Dermatology*. 2nd ed. London: Elsevier; 2008. p. 1421-35.
 72. Miyara M, Amoura Z, Parizot C, Badoual C, Dorgham K, Trad S, et al. The immune paradox of sarcoidosis and regulatory T cells. *J Exp Med* 2006;203:359-70.
 73. Cameron AL, Kirby B, Griffiths CE. Circulating natural killer cells in psoriasis. *Br J Dermatol* 2003;149:160-4.
 74. Kaštelan M, Prpić Massari L, Gruber F, Zamolo G, Zauhar G, Coklo M, et al. Perforin expression is upregulated in the epidermis of psoriatic lesions. *Br J Dermatol* 2004;151:831-6.
 75. Messenger AG, De Berker DA, Sinclair RD. Disorders of hair. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 66.1-66.100.
 76. Randall VA, Hibberts NA, Thornton MJ, Hamada K, Merrick AE, Kato S, et al. The hair follicle: A paradoxical androgen target organ. *Horm Res* 2000;54:243-50.
 77. Thornton MJ, Hamada K, Messenger AG, Randall VA. Androgen-dependent beard dermal papilla cells secrete autocrine growth factor (s) in response to testosterone unlike scalp cells. *J Invest Dermatol* 1998;111:727-32.
 78. Inui S, Fukuzato Y, Nakajima T, Yoshikawa K, Itami S. Androgen-inducible TGF-beta1 from balding dermal papilla cells inhibits epithelial cell growth: A clue to understand paradoxical effects of androgen on human hair growth. *FASEB J* 2002;16:1967-9.
 79. James WD, Berger TG, Elston DM. Atopic dermatitis, eczema, and noninfectious immunodeficiency disorders. In: James WD, Berger TG, Elston DM, editors. *Andrews' Diseases of the Skin Clinical Dermatology*. 11th ed. Philadelphia: Elsevier Saunders; 2011. p. 62-87.
 80. Staples ER, McDermott EM, Reiman A, Byrd PJ, Ritchie S, Taylor AM, et al. Immunodeficiency in ataxia telangiectasia is correlated strongly with the presence of two null mutations in the ataxia telangiectasia mutated gene. *Clin Exp Immunol* 2008;153:214-20.
 81. Weatherhead S, Robson SC, Reynolds NJ. Management of psoriasis in pregnancy. *BMJ* 2007;334:1218-20.
 82. Murase JE, Chan KK, Garite TJ, Cooper DM, Weinstein MD. Hormonal effect on psoriasis in pregnancy and post partum. *Arch Dermatol* 2005;141:601-6.
 83. Boyd AS, Morris LF, Phillips CM, Menter MA. Psoriasis and pregnancy: Hormone and immune system interaction. *Int J Dermatol* 1996;35:169-72.
 84. McNeill ME. Multiple pregnancy-induced remissions of psoriatic arthritis: Case report. *Am J Obstet Gynecol* 1988;159:896-7.
 85. Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, et al. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006;4:295-306.
 86. de Laat B, Wu XX, van Lummel M, Derksen RH, de Groot PG, Rand JH. Correlation between antiphospholipid antibodies that recognize domain I of beta2-glycoprotein I and a reduction in the anticoagulant activity of annexin A5. *Blood* 2007;109:1490-4.
 87. Horstman LL, Jy W, Bidot CJ, Ahn YS, Kelley RE, Zivadinov R, et al. Antiphospholipid antibodies: Paradigm in transition. *J Neuroinflammation* 2009;6:3.
 88. James WD, Berger TG, Elston DM. Atopic dermatitis, eczema, and

- noninfectious immunodeficiency disorders. In: James WD, Berger TG, Elston DM, editors. *Andrews' Diseases of the Skin Clinical Dermatology*. 11th ed. Philadelphia: Elsevier Saunders; 2011. p. 81-97.
89. Crowson AN. Superficial and deep perivascular dermatitis. In: Barnhill RL, Crowson AN, Magro CM, Pipekorn MW, editors. *Dermatopathology*. 3rd ed. New York: McGraw Hill companies, Inc; 2010. p. 421-45.
90. Su W, Hall BJ 4th, Cockerell CJ. Photodermatitis with minimal inflammatory infiltrate: Clinical inflammatory conditions with discordant histologic findings. *Am J Dermatopathol* 2006;28:482-5.
91. Yaar M, Gilchrist BA. Aging of skin. In: Wolff K, Goldsmith LA, Katz SI, Gilchrist BA, Paller AS, Leffell DJ I, editors. *Fitzpatrick's Dermatology in General Medicine*. 7th ed. New York: The McGraw-Hill Companies; 2008. p. 963-73.
92. Mancini G, Orange AP, den Hollander JC, Levy ML. Fibromatoses, hyalinoses and stiff skin syndrome. In: Irvine A, Hoeger P, Yan A, editors. *Harper's Textbook of Pediatric Dermatology*. 3rd ed. Oxford: Wiley-Blackwell; 2011. p. 97.1-70.21.
93. Whittaker SJ. Cutaneous lymphomas and lymphocytic infiltrates. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 57.1-57.64.
94. Pritchard BN, Youngberg GA. Atypical mitotic figures in basal cell carcinoma. A review of 208 cases. *Am J Dermatopathol* 1993;15:549-52.
95. Bowen AR, Hanks AN, Murphy KJ, Florell SR, Grossman D. Proliferation, apoptosis, and survivin expression in keratinocytic neoplasms and hyperplasias. *Am J Dermatopathol* 2004;26:177-81.
96. Miyachi Y, Ishikawa O. Dermal connective tissue metabolism in photoageing. *Australas J Dermatol* 1998;39:19-23.
97. Seite S, Zucchi H, Septier D, Igondjo-Tchen S, Senni K, Godeau G. Elastin changes during chronological and photo-ageing: The important role of lysozyme. *J Eur Acad Dermatol Venereol* 2006;20:980-7.
98. Battie C, Verschoore M. Cutaneous solar ultraviolet exposure and clinical aspects of photodamage. *Indian J Dermatol Venereol Leprol* 2012;78:S9-S14.
99. Lee SW, Byun JY, Choi YW, Myung KB, Choi HY. Fexofenadine-induced urticaria. *Ann Dermatol* 2011;23(Suppl 3):S329-32.
100. Glenn CJ, Kobraei KB, Russo JJ. New-onset psoriasis associated with adalimumab: A report of two cases. *Dermatol Online J* 2011;17:15.
101. Shelling ML, Vitiello M, Lanuti EL, Miteva M, Romanelli P, Kerdel FA. A case of palmoplantar pustulosis Induced by certolizumab pegol: New anti-TNF-alpha demonstrates the same class effect. *J Clin Aesthet Dermatol* 2012;5:40-1.
102. de Gannes GC, Ghoreishi M, Pope J, Russell A, Bell D, Adams S, *et al.* Psoriasis and pustular dermatitis triggered by TNF- α inhibitors in patients with rheumatologic conditions. *Arch Dermatol* 2007;143:223-31.
103. Satriano RA, Abbate G, Esposito S, Cassaglia B, Piccolo V, Baroni A. "Paradoxical" adverse effects caused by anti-tumor necrosis factor-alpha biological drugs: Appearance of psoriasis in a patient treated with infliximab for rheumatoid arthritis. *Indian J Dermatol Venereol Leprol* 2011;77:536.
104. Collamer AN, Guerrero KT, Henning JS, Battafarano DF. Psoriatic skin lesions induced by tumor necrosis factor antagonist therapy: A literature review and potential mechanisms of action. *Arthritis Rheum* 2008;59:996-1001.
105. Aeberli D, Seitz M, Juni P, Villiger PM. Increase of peripheral CXCR3 positive T lymphocytes upon treatment of RA patients with TNF- α inhibitors. *Rheumatology (Oxford)* 2005;44:172-5.
106. Bartolucci P, Ramanoelina J, Cohen P, Mahr A, Godmer P, Le Hello C, *et al.* Efficacy of the anti-TNF-alpha antibody infliximab against refractory systemic vasculitides: An open pilot study on 10 patients. *Rheumatology (Oxford)* 2002;41:1126-32.
107. Sokumbi O, Wetter DA, Makol A, Warrington KJ. Vasculitis associated with tumor necrosis factor- α inhibitors. *Mayo Clin Proc* 2012;87:739-45.
108. McCain ME, Quinet RJ, Davis WE. Etanercept and infliximab associated with cutaneous vasculitis. *Rheumatology (Oxford)* 2002;41:116-7.
109. Moustou AE, Matekovits A, Dessinioti C, Antoniou C, Sfrikakis PP, Stratigos AJ. Cutaneous side effects of anti-tumor necrosis factor biologic therapy: A clinical review. *J Am Acad Dermatol* 2009;61:486-504.
110. Klein R, Rosenbach M, Kim EJ, Kim B, Werth VP, Dunham J. Tumor necrosis factor inhibitor-associated dermatomyositis. *Arch Dermatol* 2010;146:780-4.
111. Daïen CI, Monnier A, Claudepierre P, Constantin A, Eschard JP, Houvenagel E, *et al.* Sarcoid-like granulomatosis in patients treated with tumor necrosis factor blockers: 10 cases. *Rheumatology (Oxford)* 2009;48:883-6.
112. Layton AM. Disorders of the sebaceous glands. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 42.1-42.89.
113. Pereira MF, Roncada EM, Oliveira CM, Monteiro R, Abreu MA, Ortigosa LC. Acne fulminans and isotretinoin: Case report. *An Bras Dermatol* 2011;86:983-5.
114. Chivot M. Acne flare-up and deterioration with oral isotretinoin. *Ann Dermatol Venereol* 2001;128 (3 Pt 1):224-8.
115. Romiti R, Jansen T, Plewig G. Acne fulminans. *An Bras Dermatol* 2000;75:611-7.
116. Berth-Jones J. Rosacea, perioral dermatitis and similar dermatoses, flushing and flushing syndromes. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. Oxford: Wiley-Blackwell; 2010. p. 43.1-43.20.
117. Fujiwara S, Okubo Y, Irisawa R, Tsuboi R. Rosaceiform dermatitis associated with topical tacrolimus treatment. *J Am Acad Dermatol* 2010;62:1050-2.
118. Antille C, Saurat JH, Lübke J. Induction of rosaceiform dermatitis during treatment of facial inflammatory dermatoses with tacrolimus ointment. *Arch Dermatol* 2004;140:457-60.
119. El Sayed F, Ammoury A, Dhaybi R, Bazex J. Rosaceiform eruption to pimecrolimus. *J Am Acad Dermatol* 2006;4:548-50.
120. Adnan ZA. Diagnosis and treatment of scleroderma. *Acta Med Indones* 2008;40:109-12.
121. Bernstein RM, Hall MA, Gostelow BE. Morphea-like reaction to D-penicillamine therapy. *Ann Rheum Dis* 1981;40:42-4.
122. Liddle BJ. Development of morphea in rheumatoid arthritis treated with penicillamine. *Ann Rheum Dis* 1989;48:963-4.
123. Zolaly MA. Histamine H1 antagonists and clinical characteristics of febrile seizures. *Int J Gen Med* 2012;5:277-81.
124. Yamashita Y, Isagai T, Seki Y, Ohya T, Nagamitsu S, Matsuishi T. West syndrome associated with administration of a histamine H1 antagonist, oxatomide. *Kurume Med J* 2004;51:273-5.
125. Dierickx CC. Laser treatment of pigmented lesions. In: Goldberg DJ, editor. *Laser Dermatology*. 1st ed. Berlin: Springer; 2005. p. 36-60.
126. Alajlan A, Shapiro J, Rivers JK, MacDonald N, Wiggin J, Lui H. Paradoxical hypertrichosis after laser epilation. *J Am Acad Dermatol* 2005;53:85-8.

Cite this article as: Adya KA, Inamadar AC, Palit A. Paradoxes in dermatology. *Indian Dermatol Online J* 2013;4:133-42.

Source of Support: Nil, **Conflict of Interest:** None declared.