

Comparison of Three Blood Culture Techniques in the Diagnosis of Human Brucellosis

Smita Mangalgi, Annapurna Sajjan

Department of Microbiology, BLDEU's Shri B. M. Patil Medical College, Bijapur, Karnataka, India

Address for correspondence: Dr. Smita Mangalgi, E-mail: smitamangalgi@gmail.com

ABSTRACT

Background: Isolation of *Brucella* is the gold standard in the laboratory diagnosis of brucellosis. As *Brucella* is intracellular and the number of circulating bacteria is usually low, removal/dilution of antibacterial substances, concentration of bacteria and optimal culture conditions may enhance the rate of isolation.

Aims and Objectives: The objective of the following study was to compare the lysis concentration (LC), clot culture and conventional Castaneda blood culture techniques for the isolation rate and recovery time in the diagnosis of human brucellosis.

Materials and Methods: Blood cultures by LC, clot culture and conventional method were performed in 169 patients who had antibody titers ≥ 160 international units by the serum agglutination test.

Results: Overall blood culture positivity was found to be 24.8%, 43.1% and 34.9% by conventional, LC and clot culture techniques in that order. The mean recovery time by LC and clot culture techniques was significantly less than conventional method, resulting in an overall difference of nearly 6 and 4 days respectively.

Conclusions: For the isolation of *Brucella* from blood, LC and clot culture techniques are better than the conventional technique.

Key words: Blood culture, *brucella*, clot culture, conventional, lysis concentration

INTRODUCTION

Brucellosis is endemic in low socio-economic countries like India and the prevalent species causing human infections are *Brucella melitensis* and *Brucella abortus*.^[1] Contact with the infected animals, consumption of raw milk and ignorance regarding the disease, are the major risk factors.^[1,2] Since clinical manifestations of this disease are protean in nature, laboratory help is a must in the diagnosis. In the laboratory, brucellosis is diagnosed either by isolation of the causative agent and/or by serological tests. Although an array of serological tests is available, Rose Bengal Plate Test (RBPT) and serum agglutination test (SAT) are the most widely used. The sensitivity

of RBPT is considerably high, but its specificity is much lower when used to test individuals residing in an endemic area.^[3] In the same way, SAT, used to confirm RBPT results, also has limitations of lack of sensitivity as well as specificity.^[4-7] Apart from this, serological testing does not provide direct evidence for the presence of the pathogen, hence, isolation of *Brucella* spp. from the clinical specimen is considered to be the gold standard.^[2,8,9]

Many of the methods have been developed for culturing *Brucella* spp. from blood specimen, including the conventional Ruiz-Castaneda (Castaneda) method, automated systems, lysis concentration (LC) and clot culture.^[10-14] In general, blood culture for *Brucella* is performed by conventional Castaneda method where the blood specimen is directly inoculated in the liquid phase of the Castaneda medium. Although the results by this method are satisfactory in acute untreated cases, the incubation time required is very long. The isolation rate is markedly reduced in treated cases or sub-acute/chronic brucellosis patients.^[13] The earlier reports have shown an increase in isolation rate and

Access this article online

Quick Response Code:



Website:
www.jlponline.org

DOI:
10.4103/0974-2727.129084

decrease in recovery time by adapting LC technique.^[12,13] However, both conventional and LC methods require one more blood sample for culture once the serological test is found positive. As it is difficult to obtain one more blood specimen in ignorant patients, clot culture could be a better option. Isolation of *Salmonella* by clot culture has been tried by many workers.^[15-17] Reports on clot culture for *Brucella* are scarce and controversial.^[15,18] Hence in this study, an attempt has been made to compare LC, clot culture and conventional culture techniques for the isolation of *Brucella* from blood specimen in an endemic area.

MATERIALS AND METHODS

Blood culture by all the three techniques was performed in 169 out of 191 patients showing SAT titer of ≥ 160 international units (IU) after obtaining written consent. Depending on the duration of the disease, patients were divided into three groups as: acute (<8 weeks), sub-acute (8-52 weeks) and chronic (>52 weeks) brucellosis.^[15,18] Patients were considered to have relapse/re-infection if symptoms reappeared within a year after the completion of antibrucellar treatment.

Five ml of blood was collected aseptically from patients with the clinical symptoms resembling brucellosis (fever, malaise, joint pain, low backache, headache, and weight loss) in a sterile, plastic screw capped bottle with 6-7 glass beads of 2 mm diameter. The sample was allowed to clot. The specimen was centrifuged at 2000 rpm for 15 min; the serum was separated aseptically and used for serology. The clot was preserved in the refrigerator for clot culture if needed.

Blood sample for conventional and LC culture technique was collected from patients who had significant SAT titers (≥ 160 IU). About 10 ml of blood was collected in adults and 5 ml in case of children. The specimen was equally distributed and processed for conventional blood culture and LC technique.

Conventional culture technique

The blood specimen was inoculated aseptically into the broth phase of Castaneda's biphasic medium consisting of brain heart infusion agar and broth with *Brucella* selective supplement (Hi-Media).

LC technique

Procedure described by Etemadi *et al.*, was employed with a minor modification.^[12] 5 ml of blood was added to 40 ml

of sterile distilled water with 1.5 ml of 4% sodium citrate in adults and 2.5 ml was added to 20 ml of sterile distilled water with 0.75 ml of 4% sodium citrate in children, in a sterile screw capped centrifuge tube. The contents were gently mixed and the tubes were centrifuged at 7,000 rpm for 30 min. The supernatant was discarded and the sediment was inoculated in the Castaneda's medium, instead of culture plate.

Clot culture technique

The blood clot preserved in the sterile screw capped plastic tube with glass beads after removal of serum was used for this method. The clot was disrupted by shaking the tubes on a shaker for 15 min. The disrupted clot was then inoculated in Castaneda's medium.

Incubation

The media were incubated at 37°C with 10% CO₂ for a maximum of 30 days.

Recording of results

The bottles were observed daily for growth. Sub-culturing was done by allowing the blood broth mixture to flow over the solid phase. The day of appearance of the first colony was recorded for comparison of growth rates.

Identification and confirmation of the isolates

The isolates were identified based on colony morphology, gram stain, modified "ZN" staining, CO₂ requirement, biochemical tests like oxidase and urease, H₂S production for 4 days and growth in the presence of basic fuchsin (1:50,000 and 1:100,000) and thionin (1:25,000, 1:50,000 and 1:100,000). Provisional confirmation and bio-typing of the isolate was done by performing slide agglutination test using *B. abortus* and *B. melitensis* monospecific antisera (Murex Biotech). The isolates were confirmed at IVRI Izatnagar/DRDE Gwalior.

The study was approved by the Institutional Ethical Committee.

Statistical analysis

Statistical analysis of the data was performed by GraphPad InStat software designed by GraphPad Software, Inc. USA. Chi-square test with Yate's correction was used for comparison.

RESULTS

Blood culture by all the three techniques was performed in 169 out of 191 patients. Overall blood culture positivity was found to be 24.8%, 43.1% and 34.9% by conventional, LC and clot culture techniques respectively. The isolation rates in patients with acute, sub-acute, chronic brucellosis and relapse/reinfection by various culture techniques are given in Table 1.

When compared with the conventional technique LC and clot culture techniques yielded growth in significantly more number of cases ($P < 0.0001$). Isolation rate was high in acute stage compared with sub-acute stage. *Brucella* could not be isolated in chronic patients by any of the methods.

The mean recovery time for conventional technique was 9.61 days (standard deviations [SD] ± 1.68), whereas with LC and clot culture techniques it was 4.08 (SD ± 0.87 ; $P < 0.0001$) and 5.83 (SD ± 1.37 ; $P < 0.0001$) respectively. In the acute stage, recovery time was relatively less by all the three methods [Table 2]. Earliest growth by the conventional technique was seen on the 6th day where as it was on the 3rd day with both LC and clot culture techniques. LC technique showed earlier growth in 96% blood cultures, before the first positive culture by conventional method was noted, whereas clot culture yielded earlier growth in 42.55% of cultures. Maximum incubation time required for culture to be positive by LC, clot culture and conventional technique was 6, 9 and 16 days respectively. After 16th day, no culture was found positive, though the cultures were incubated for 1 month. Despite maximum aseptic precautions taken during collection and processing of the sample, contamination rate was high for clot culture (5.32%) followed by LC (2.95%), may be due to more number of manipulations required for the procedure. None of the conventional cultures showed contamination.

DISCUSSION

In the present study, blood culture positivity was found to be 24.8%, 43.1% and 34.9% and the recovery time was 10, 4 and 6 days by conventional, LC and clot culture techniques respectively. Our findings suggest that LC and clot culture methods are more sensitive and require less time than conventional culture thereby helping in early detection of infection. Among the LC and clot culture techniques, LC was better due to increased isolation rates and decreased recovery time. Similar findings have been reported by Etemadi *et al.*, Mantur and Mangalgi and Espinosa *et al.*,^[12-14]

The mean recovery time by LC in this study was 4.08 days, which is in agreement with Espinosa *et al.*,^[14] but slightly more as compared to the results of 2 and 2.7 days by Etemadi *et al.*, and Mantur and Mangalgi.^[12,13]

In a study done by Escamilla *et al.*, clot culture technique was found to be less sensitive and more labor-intensive when compared with the conventional method.^[15] Conversely in our study, clot culture technique was more sensitive and also the recovery time was less when compared with the conventional method. Similar findings have been reported by Mantur *et al.*,^[18] Another advantage of this technique is that need for one more blood sample is alleviated.

In a study by Espinosa *et al.*, isolation rate was higher in sub-acute cases than in patients with acute brucellosis by LC technique.^[14] Whereas far higher isolation rates in both acute and chronic stages have been reported by Mantur and Mangalgi.^[13] In our study, culture positivity rate was high in patients with acute followed by sub-acute brucellosis and none of the chronic cases grew *Brucella*.

As the patients in our study were not relieved of their symptoms over a long time, they switched on to many doctors before being diagnosed as brucellosis cases at our hospital, which could be an attribute for our lower isolation rates. History of minimum 3-4 courses of antibiotics was elicited in 116 (68.63%) patients.

In general, blood cultures are attempted to confirm the presumptive serological diagnosis. In India, especially in rural areas, due to the misperception that blood once lost is lost

Table 1: Culture positivity by various culture techniques in different stages of brucellosis

Disease stage	No. of patients with SAT titer ≥ 160 IU	Culture done	Number of specimens positive by (%)		
			Conventional	Lysis	Clot
Acute	125	103	37 (35.9)	58 (56.3)***	47 (45.6)***
Sub-acute	35	35	3 (8.5)	13 (37.1)**	10 (28.5)**
Chronic	29	29	0	0	0
Relapse/reinfection	2	2	2 (100)	2 (100)*	2 (100)*
Total	191	169	42 (24.8)	73 (43.9)***	59 (34.91)***

SAT: Serum agglutination test, IU: International units. ***Extremely significant, **Significant, *Not significant

Table 2: Recovery time in days by different culture methods in various stages of brucellosis

Disease stage	Culture method		
	Conventional	Lysis	Clot
Acute	9.45 \pm 1.6	4 \pm 0.85	5.63 \pm 1.3
Sub-acute	12.75 \pm 2.75	4.61 \pm 0.76	6.80 \pm 1.47
Relapse/reinfection	9.5 \pm 0.7	3 \pm 0.00	5.5 \pm 0.70
Overall	9.61 \pm 1.68	4.08 \pm 0.87	5.83 \pm 1.37

Mangalgi and Sajjan: Blood culture in human brucellosis

forever and blood loss enhances the disease; it is difficult to obtain one more blood specimen for culture in seropositive individuals. In such situations and also in field conditions (camps), wherein obtaining a second sample for culture is difficult, clot culture could be the optimal alternative. Both lysis and clot culture require manipulations, which enhance the chances of laboratory infection and contamination, thereby necessitating strict biosafety and aseptic measures.

CONCLUSIONS

We propose that for the isolation of *Brucella* from blood specimen, LC method is better than conventional Castaneda's method as the isolation rate is high and the recovery time is less. Clot culture is a better option when a second blood sample cannot be obtained for culture. As lysis and clot culture techniques are sensitive, simple and inexpensive and yield earlier results, they can be adapted in the technically and economically backward areas where automated systems are far from reach. However, to prevent laboratory acquired infection, these methods are to be carried out in Class II biosafety cabinet with BSL-3 precautions.

REFERENCES

1. Mantur BG, Amarnath SK. Brucellosis in India-A review. *J Biosci* 2008;33:539-47.
2. Corbel MJ. Brucellosis in Humans and Animals. WHO Press, Geneva, Switzerland; 2006.
3. Serra J, Viñas M. Laboratory diagnosis of brucellosis in a rural endemic area in Northeastern Spain. *Int Microbiol* 2004;7:53-8.
4. Gómez MC, Nieto JA, Rosa C, Geijo P, Escribano MA, Muñoz A, *et al.*

- Evaluation of seven tests for diagnosis of human brucellosis in an area where the disease is endemic. *Clin Vaccine Immunol* 2008;15:1031-3.
5. Vancelik S, Guraksin A, Ayyildiz A. Seroprevalence of human brucellosis in rural endemic areas in eastern Turkey. *Trop Doct* 2008;38:42-3.
6. Agasthya AS, Isloor S, Krishnamsetty P. Seroprevalence study of human brucellosis by conventional tests and indigenous indirect enzyme-linked immunosorbent assay. *ScientificWorldJournal* 2012;2012:104239.
7. Clavijo E, Díaz R, Anguita A, García A, Pinedo A, Smits HL. Comparison of a dipstick assay for detection of *Brucella*-specific immunoglobulin M antibodies with other tests for serodiagnosis of human brucellosis. *Clin Diagn Lab Immunol* 2003;10:612-5.
8. Nielsen K, Yu WL. Serological diagnosis of brucellosis. *Sec Biol Med Sci* 2010; XXXI/1:65-89.
9. Pappas G, Akritidis N, Bosilkovski M, Tsianos E. Brucellosis. *N Engl J Med* 2005;352:2325-36.
10. Ruiz J, Lorente I, Pérez J, Simarro E, Martínez-Campos L. Diagnosis of brucellosis by using blood cultures. *J Clin Microbiol* 1997;35:2417-8.
11. Yagupsky P. Detection of *Brucellae* in blood cultures. *J Clin Microbiol* 1999; 37:3437-42.
12. Etemadi H, Raissadat A, Pickett MJ, Zafari Y, Vahedifar P. Isolation of *Brucella* spp. from clinical specimens. *J Clin Microbiol* 1984;20:586.
13. Mantur BG, Mangalgi SS. Evaluation of conventional castaneda and lysis centrifugation blood culture techniques for diagnosis of human brucellosis. *J Clin Microbiol* 2004;42:4327-8.
14. Espinosa BJ, Chacaltana J, Mulder M, Franco MP, Blazes DL, Gilman RH, *et al.* Comparison of culture techniques at different stages of brucellosis. *Am J Trop Med Hyg* 2009;80:625-7.
15. Escamilla J, Florez-Ugarte H, Kilpatrick ME. Evaluation of blood clot cultures for isolation of *Salmonella typhi*, *Salmonella paratyphi-A*, and *Brucella melitensis*. *J Clin Microbiol* 1986;24:388-90.
16. Abhijit A, Sunita N. The study of salmonellosis with reference to salmonella typhi in enteric fever patients. *Journal of Clinical and Diagnostic Research*. 2011;5:467-9.
17. Peerapur BV, Smitha S. Comparison of whole blood culture and blood clot culture for the diagnosis of enteric fever. *JKIMSU* 2013;2:145-6.
18. Mantur BG, Bidari LH, Akki AS, Mulimani MS, Tikare NV. Diagnostic yield of blood clot culture in the accurate diagnosis of enteric fever and human brucellosis. *Clin Lab* 2007;53:57-61.

How to cite this article: Mangalgi S, Sajjan A. Comparison of three blood culture techniques in the diagnosis of human Brucellosis. *J Lab Physicians* 2014;6:14-7.

Source of Support: Nil. **Conflict of Interest:** None declared.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) First Page File:

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) Article File:

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1024 kb. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) Images:

Submit good quality color images. Each image should be less than **4096 kb (4 MB)** in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) Legends:

Legends for the figures/images should be included at the end of the article file.